

Actinic keratosis

Author: Dr Amanda Oakley, Dermatologist, Hamilton, New Zealand, 1997. Updated December 2015.

What is an actinic keratosis?

Actinic keratosis is a precancerous scaly spot found on sun-damaged skin, also known as solar keratosis. It may be considered an early form of cutaneous [squamous cell carcinoma](#) (a [keratinocyte cancer](#)).

Actinic keratoses



Who gets actinic keratoses?

Actinic keratoses affect people that have often lived in the tropics or subtropics and have predisposing factors such as:

- Other signs of photoageing skin
- Fair skin with a history of sunburn
- History of long hours spent outdoors for work or recreation
- Defective immune system.

What causes actinic keratoses?

Actinic keratoses are the result of abnormal skin cell development due to DNA damage by short wavelength UVB.

They are more likely to appear if the immune function is poor, due to ageing, recent sun exposure, [predisposing disease](#), or certain [drugs](#).

What are the clinical features of actinic keratosis?

Actinic keratosis may be solitary but there are often multiple keratoses. The appearance varies.

- A flat or thickened papule, or plaque
- White or yellow; scaly, warty, or horny surface
- Skin coloured, red, or pigmented
- Tender or asymptomatic

Actinic keratoses are very common on sites repeatedly exposed to the sun, especially the backs of the hands and the face, most often affecting the ears, nose, cheeks, upper lip, vermilion of the lower lip, temples, forehead, and balding scalp. In severely chronically sun-damaged individuals, they may also be found on the upper trunk, upper and lower limbs, and dorsum of feet.

Actinic keratoses



Many more images of actinic keratoses ...

- Keratoses on the nose
- Keratoses on the face
- Keratoses on the scalp
- Keratoses on the hands
- Keratoses on the legs
- Keratoses treated with imiquimod

Complications of actinic keratoses

The main concern is that actinic keratoses indicate an increased risk of developing cutaneous [squamous cell carcinoma](#). It is rare for a solitary actinic keratosis to evolve to squamous cell carcinoma (SCC), but the risk of SCC occurring at some stage in a patient with more than 10 actinic keratoses is thought to be about 10 to 15%. A tender, thickened, ulcerated, or enlarging actinic keratosis is suspicious of evolution to SCC.

[Cutaneous horn](#) may arise from an underlying actinic keratosis or SCC.

Because they are sun damaged, people with actinic keratoses are also at risk of developing [actinic cheilitis](#), [basal cell carcinoma](#) (BCC, which is more common than SCC), [melanoma](#), and rare forms of skin cancer such as [Merkel cell carcinoma](#).

How is an actinic keratosis diagnosed?

Actinic keratosis is usually easy to diagnose clinically or by [dermoscopy](#) (see [actinic keratosis dermoscopy](#)). Occasionally, a [biopsy](#) is necessary, for example, to exclude [SCC](#), or if treatment fails.

What is the treatment for actinic keratoses?

Actinic keratoses are usually removed because they are unsightly or uncomfortable, or because of the risk that skin cancer may develop in them.

Treatment of an actinic keratosis requires removal of the defective skin cells. Epidermis regenerates from surrounding or follicular keratinocytes that have escaped sun damage.

Tender, thickened, ulcerated, or enlarging actinic keratoses should be treated aggressively. Asymptomatic flat keratoses may not require active treatment but should be kept under observation.

Physical treatments

Physical treatments are used to destroy individual keratoses that are generally symptomatic or have a thick hard surface scale. The lesions may recur in time, in which case they may be retreated by the same or a different method.

Cryotherapy using liquid nitrogen

[Liquid nitrogen](#) spray is required to achieve adequate depth and duration of the freeze. This varies according to lesion location, width, and thickness. Healing varies from 5–10 days on the face, 3–4 weeks on the hands, and 6 weeks or longer on the l

Privacy

freeze for a superficial actinic keratosis usually leaves no mark, but longer freeze times result in hypopigmentation or scar.

Shave, curettage and electrocautery

Shave, [curettage](#) (scraping with a sharp instrument) and [electrocautery](#) (burning) may be necessary to remove a [cutaneous horn](#) or hypertrophic actinic keratosis. Healing of the wound takes several weeks or longer, depending on the body site. The specimen is sent for histological examination.

Excision

[Excision](#) ensures the actinic keratosis has been completely removed, which should be confirmed by pathology. The surgical wound is sutured (stitched). The sutures are removed after a few days, the time depending on the size and location of the lesion. The procedure leaves a permanent scar.

Field treatments

Creams are used to treat areas of sun damage and flat actinic keratoses, sometimes after physical treatments have been carried out. Field treatments are most effective on facial skin. Pretreatment with keratolytics (such as [urea](#) cream, [salicylic acid](#) ointment or [topical retinoid](#)), and thorough skin cleansing improves response rates. Results are variable and the course of treatment may need repeating from time to time. With the exception of diclofenac gel, field treatments all result in local inflammatory reactions such as redness, blistering and discomfort for a varying length of time.

- [Diclofenac](#) is more often used as an anti-inflammatory drug. Applied as a gel twice daily for 3 months, it is fairly well tolerated in the treatment of actinic keratoses, but less effective than the other options listed here.
- [5-Fluorouracil](#) is a cytotoxic agent. The cream formulation is applied once or twice daily for 2 to 8 weeks. 5-fluorouracil cream is sometimes combined with [salicylic acid](#). Its effect may be enhanced by [calcipotriol](#) ointment.
- [Imiquimod](#) cream is an immune response modifier. It is applied 2 or 3 times weekly for 4 to 16 weeks.
- [Photodynamic therapy \(PDT\)](#) involves applying a photosensitiser (a porphyrin chemical such as [methyl aminolevulinic acid](#)) to the affected area prior to exposing it to a source of visible light.
- [Ingenol mebutate gel](#) is effective after only 3 applications.

Prevention of actinic keratoses

Actinic keratoses are prevented by strict [sun protection](#). If already present, actinic keratoses may improve with a very high sun protection factor (50+) broad-spectrum [sunscreen](#) applied at least daily to affected areas, year-round.

The number and severity of actinic keratoses can also be reduced by taking [nicotinamide](#) (vitamin B3) 500 mg twice daily.

What is the outlook for actinic keratoses?

Actinic keratoses may recur months or years after treatment. The same treatment can be repeated or another method used. Patients who have been treated for actinic keratoses are at risk of developing new keratoses. They are also at increased risk of other [skin cancers](#), especially [intraepidermal squamous cell carcinoma](#), invasive cutaneous [squamous cell carcinoma](#), [basal cell carcinoma](#) and [melanoma](#).